

# Indiana State Department of Health

## Minimum Immunization Requirements for School Entry 2011-2012

### **Pre-Kindergarten**

- 4 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) or any combination of the three are required.
- 3 doses of either oral polio (OPV) or inactivated polio (IPV) vaccine in any combination
- 3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).
- 1 doses of measles (rubeola) vaccine, on or after the first birthday.
- 1 doses of mumps vaccine, on or after the first birthday.
- 1 dose of rubella (German measles) vaccine, on or after the first birthday.
- **2 doses of varicella (chickenpox) vaccine on or after the first birthday and separated by 3 months or physician written documentation of history of chicken pox disease, including month and year of disease.**

### **Kindergarten**

- 5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) or (4 doses are acceptable if the 4<sup>th</sup> dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).
- 4 doses of any combination of IPV or OPV. **The 4<sup>th</sup> dose must be administered on or after the 4<sup>th</sup> birthday, and at least 6 months after the previous dose. (3 doses of all OPV or all IPV are acceptable if the 3<sup>rd</sup> dose was administered on or after the 4<sup>th</sup> birthday, and at least 6 months after the 2<sup>nd</sup> dose).**
- 3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be given on or after 24 weeks of age and no earlier than 16 weeks after the first birthday).
- 2 doses of measles (rubeola) vaccine, on or after the first birthday.
- 2 doses of mumps vaccine, on or after the first birthday.
- 1 dose of rubella (German measles) vaccine, on or after the first birthday.
- **2 doses of varicella (chickenpox) vaccine on or after the first birthday and separated by 3 months or physician written documentation of history of chicken pox disease, including month and year of disease.**

## **Grade 1**

- 5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).
- 4 doses of any combination of IPV or OPV by age 4-6, (3 doses of all OPV or IPV are acceptable if the 3rd dose was administered on or after the 4th birthday).
- 3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).
- 2 doses of measles (rubeola) vaccine, on or after the first birthday.
- 2 doses of mumps vaccine, on or after the first birthday.
- 1 dose of rubella (German measles) vaccine, on or after the first birthday.
- 1 dose of varicella (chickenpox) vaccine on or after the first birthday **or physician written documentation of history of chicken pox disease, including month and year of disease.**

## **Grades 2-5**

- 5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).
- 4 doses of any combination of IPV or OPV by age 4-6, (3 doses of all OPV or IPV are acceptable if the 3rd dose was administered on or after the 4th birthday).
- 3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).
- 2 doses of measles (rubeola) vaccine, on or after the first birthday.
- 2 doses of mumps vaccine, on or after the first birthday.
- 1 dose of rubella (German measles) vaccine, on or after the first birthday.
- 1 dose of varicella (chickenpox) vaccine on or after the first birthday or written history of disease. Parental history of chickenpox disease is acceptable proof of immunity. A signed written statement from the parent/guardian indicating month and year of disease is sufficient.

## **Grades 6-12**

- 5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).
- 4 doses of any combination of IPV or OPV by age 4-6, (3 doses of all OPV or IPV are acceptable if the 3rd dose was administered on or after the 4th birthday).
- 3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on after after 24 weeks of age).
- 2 doses of measles (rubeola) vaccine, on or after the first birthday.
- 2 doses of mumps vaccine, on or after the first birthday.
- 1 dose of rubella (German measles) vaccine, on or after the first birthday.
- **2 doses of varicella (chickenpox) vaccine on or after the first birthday separated by age-appropriate interval or written history of disease. Parental history of chickenpox disease is acceptable proof of immunity. A signed written statement from the parent/guardian indicating month and year of disease is sufficient.**
- **1 dose of tetanus-diphtheria-acellular pertussis vaccine (Tdap) given on or after 10 years of age.**
- **1 dose of meningococcal conjugate vaccine (MCV4).**

\* For children who have delayed immunizations, please refer to the 2010 CDC "Catch-up Immunization Schedule" to determine adequately immunizing doses. All minimum intervals and ages for each vaccination as specified per 2010 CDC guidelines must be met for a dose to be valid. These guidelines can be found a [www.cdc.gov/vaccines/recs/schedules/default.htm](http://www.cdc.gov/vaccines/recs/schedules/default.htm)